

Received Date : 16-Dec-2013  
Accepted Date : 02-Apr-2014  
Article type : Main Research Article

## **Prevalence of emotional, physical and sexual abuse among pregnant women in six European countries**

Running title: Abuse among European pregnant women

Mirjam LUKASSE<sup>1,2</sup>, Anne-Mette SCHROLL<sup>3</sup>, Elsa Lena RYDING<sup>4</sup>, Jacquelyn CAMPBELL<sup>5</sup>, Helle KARRO<sup>6</sup>, Hildur KRISTJANSDOTTIR<sup>7,8</sup>, Made LAANPERE<sup>9</sup>, Thora STEINGRIMSDOTTIR<sup>10</sup>, Ann TABOR<sup>3,11</sup>, Marleen TEMMERMAN<sup>12</sup>, An-Sofie VAN PARYS<sup>12</sup>, Anne-Marie WANGEL<sup>13</sup> & Berit SCHEI<sup>1,14</sup>

<sup>1</sup> Department of Public Health and General Practice, Norwegian University of Science and Technology, Trondheim, Norway

<sup>2</sup> Faculty of Health Sciences, Oslo and Akershus University College of Applied Sciences, Department of Obstetrics and Gynecology, St.Olavs University Hospital, Trondheim, Norway.

<sup>3</sup> Center for Fetal Medicine and Pregnancy, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark

<sup>4</sup> Department of Women's and Children's Health, Division of Obstetrics and Gynecology, Karolinska Institute, Stockholm, Sweden

<sup>5</sup> Johns Hopkins University, School of Nursing, Baltimore, MD, USA

<sup>6</sup> Department of Obstetrics and Gynecology, University of Tartu, Tartu University Hospital Women's Clinic, Tartu, Estonia

<sup>7</sup> Health Directorate, Reykjavik, Iceland

<sup>8</sup> University of Iceland, Faculty of Nursing, department of Midwifery, Reykjavik, Iceland

<sup>9</sup> Department of Obstetrics and Gynecology, University of Tartu, Tartu Sexual Health Clinic, Estonia

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/aogs.12392

This article is protected by copyright. All rights reserved.

<sup>10</sup> Landspítali University Hospital, Reykjavik, Iceland

<sup>11</sup> Faculty of Health Sciences, Copenhagen University, Copenhagen Denmark

<sup>12</sup> Faculty of Medicine and Health Sciences Ghent University, International Centre for Reproductive Health, Belgium

<sup>13</sup> Malmö University, Faculty of Health and Society, Malmö, Sweden

<sup>14</sup> Department of Obstetrics and Gynecology, St.Olav's University Hospital, Trondheim, Norway

**Corresponding author:**

Mirjam Lukasse

Faculty of Health Sciences, PB 4. St. Olavs plass, 0130 Oslo, Norway

E-mail: Mirjam.Lukasse@hioa.no

**Conflict of interest**

We hereby certify that there is no conflict of interest with regarding the material discussed in the manuscript.

Mirjam Lukasse

Anne-Mette Schroll

Berit Schei

Jacquelyn Campbell

Helle Karo

Hildur Kristjansdottir

Made Laanpere

Thora Steingrimsdottir

Ann Tabor

Marleen Temmerman

An-Sofie Van Parys

Anne-Marie Wangel

Elsa Lena Ryding

## Abstract

*Objectives.* The primary objective was to investigate the prevalence of a history of abuse among women attending routine antenatal care in six Northern European countries. Secondly, we explored current suffering from reported abuse. *Design.* A prospective cohort study *Setting.* Routine antenatal care in in Belgium, Iceland, Denmark, Estonia, Norway, and Sweden between March 2008 and August 2010. *Population.* 7174 pregnant women *Methods.* A questionnaire including a validated instrument measuring emotional, physical and sexual abuse. *Main outcome measure.* Proportion of women reporting emotional, physical and sexual abuse. Severe current suffering defined as a Visual Analogue Scale score of  $\geq 6$ .

*Results.* An overall lifetime prevalence of any abuse was reported by 34.8% of the pregnant women. The ranges across the six countries of lifetime prevalence were 9.7–30.8% for physical abuse, 16.2–27.7% for emotional abuse, and 8.3–21.1% for sexual abuse. Few women reported current sexual abuse, 0.4% compared to 2.2% current physical abuse and 2.7% current emotional abuse. Current severe suffering was reported by 6.8% of the women who reported physical abuse, 9.8% of those who reported sexual abuse and 13.5% for emotional abuse. *Conclusion.* A high proportion of pregnant women attending routine antenatal care report a history of abuse. About one in ten of them experiences severe current suffering from the reported abuse. In particular these women might benefit from being identified in the antenatal care setting and being offered specialized care.

## Keywords

abuse, violence against women, pregnancy, prevalence

## List of abbreviations

NorAQ NorVold Abuse Questionnaire

OR odds ratio

G Goodman-Kruskal gamma

## Key Message

A history of abuse is common among pregnant women in Northern Europe. About one in ten women reports severe suffering from previous or current abuse. Routine antenatal care provides a window of opportunity to identify suffering and offer specialized care.

## Introduction

Abuse of women and girls is a widely recognized public health issue (1). The term abuse is generally used when violence or acts of violation are part of an ongoing pattern or behaviour. The World Health Organization defines violence against women as “any act of gender-based violence that results in, or is likely to result in physical, sexual or mental harm or suffering to women, including threats of such acts, coercion or arbitrary deprivation of liberty, whether occurring in public or private life”(1, 2).

When acts of violence and abuse occur within the privacy of the home they can also be defined as domestic violence (3), while violence inflicted by a current or previous partner falls under the term intimate partner violence (3). Partner violence is the leading cause of death among women in reproductive age (4, 5). Other detrimental consequences of abuse on mental and physical health are well documented (6-9). Evidence suggests that women are particularly vulnerable to abuse during pregnancy and the postnatal period (10, 11). Violence and abuse has been shown to influence women’s health during pregnancy and birth and may affect the health of the fetus and new-born child (12-16). The different pathways described are direct injury, neurobiological changes, and an increase in health-detrimental behaviours such as eating disorders and drug abuse (13, 15, 17).

Previously published estimates of prevalence of past and present violence and abuse among pregnant women vary greatly and may be difficult to compare, as they differ regarding the type of abuse assessed, time of occurrence, and perpetrator (11). In addition, methodological factors such as study design, measuring instrument and population studied can influence results (11). There are two previous studies presenting internationally comparable data on the population prevalence of violence against women and estimates of the occurrence during pregnancy (7, 18). These studies were restricted to intimate partner violence and thus excluded abuse women had experienced as a child and violence perpetrated by others than a present or previous intimate partner (7, 18). The method in both studies was a standardised household survey including women at all ages and asking them to recall whether violence had occurred during pregnancy (7, 18). While these studies present valuable information their relevance to a European setting is limited (7, 18). There are no international population-based studies conducted among pregnant women attending routine antenatal care, estimating the prevalence of physical, emotional and sexual violence abuse experienced as a child or/and as an adult. This was the primary aim of our study. Secondly, we explored current suffering from reported abuse.

## Material and Methods

The Bidens study, a six-country (Belgium, Iceland, Denmark, Estonia, Norway, and Sweden) cohort study of unselected pregnant women, was the result of an European Union-funded collaboration between the Norwegian University of Science and Technology (NTNU) and partners from universities and hospitals in six European countries. A short description of the study sites is given as Supporting Information Table S1. There were between one and 7 urban antenatal care sites of data collection in each country with the most in Norway (5) and Sweden (7).

Recruitment took place between March 2008 and August 2010. A total of 7200 women who consented, subsequently completed a questionnaire and allowed the extraction of specified data on their delivery from their medical notes. Due to country specific organization as well as the requirements of local ethical committees, minor variations in the recruitment procedure occurred.

In *Belgium*, women were approached by the midwife or secretary when attending antenatal care. Women were asked to complete the questionnaire in the privacy of a separate room. In *Iceland* women were recruited when attending routine ultrasound and returned completed forms by mail. In *Denmark* women were given information about the study when attending early routine ultrasound screening and were mailed the questionnaire later. They returned the questionnaire by mail or when attending their next ultrasound examination. In *Estonia* women were invited to participate while visiting for an antenatal consultation. After completing the questionnaire it was left in a mailbox at the clinic. In *Norway*, women received the questionnaire by mail and returned it by mail, after attending routine ultrasound. Non-responders were sent one reminder. In *Sweden*, the questionnaire was administered to women when attending routine glucose tolerance tests and filled out during the two hours between the blood samplings.

The right to obtain information on non-participating women varied between countries and hence the basis for calculating response rates. In *Belgium* and *Sweden* registrations of non-participants was not allowed, the response rate was estimated at 50% and 78%, respectively. In *Iceland* and *Denmark* the response rate was 65 % and 57.3 %, respectively (no reminder). In *Estonia*, the response rate was 90%, based on number of questionnaires given to the assigned study midwives and number of filled out forms returned. In *Norway* the participation rate was 50% (one reminder). The estimated response rate varied between 50% in Norway to 90% in Estonia.

All women required sufficient language skills to fill out the form. In Estonia women could choose to complete an Estonian or Russian language questionnaire. In Belgium, Iceland and Denmark women less than 18 years of age were excluded. In Denmark, only women from the local geographical area were invited. In Belgium, women who could not to be separated from their accompanying person were not recruited. In Iceland, Denmark and Norway, women with major fetal pathologies were excluded from the study.

The questionnaire included questions on socio-economic background, general and mental health and obstetric history. The questions on abuse were taken from the NorVold Abuse Questionnaire (NorAQ), which was developed in a Nordic multi-centre study among gynaecological patients (19). This validated instrument includes 13 descriptive questions measuring emotional, physical and sexual abuse (20). A complete version of the questionnaire was developed in English. Where a previously translated version of the NorAQ was available, this was used. Additional items of the NorAQ questionnaire were translated into the required languages by a native speaker (Flemish, Icelandic, Danish, Estonian, Russian, Norwegian and Swedish) and then translated back again into the source language. The original and back-translated versions were used to determine the final consensus version.

Emotional, physical and sexual abuse was assessed in three identically structured sections. For each type and level of abuse the answer categories were no, yes as a child, yes as an adult, or yes both as a child and as an adult. These were classified according to the most severe level reported (mild, moderate and, severe). Two items addressing 'mild sexual abuse with no genital contact' and 'mild humiliating sexual abuse' were combined in the analysis into one category of 'mild sexual abuse'. For each type of abuse women were asked if they experienced the indicated abuse during the past 12 months, which was coded as current. The degree of current suffering was measured on a visual analogous scale (0–10) and recoded into no suffering (0), moderate suffering (1-5) and severe current suffering ( $\geq 6$ ), based on the distribution of the data. Women were defined as having experienced any abuse if they answered yes to at least one of the questions of sexual, emotional and physical abuse. The question measuring mild physical abuse has shown low specificity in the validation study (20). Thus results are presented including and excluding this item.

The study was conducted in accordance with the ethical guidelines developed by the World Health Organization (25), which highlight the importance of ensuring women's safety,

This article is protected by copyright. All rights reserved.

confidentiality and privacy. The information letter instructed women to complete the form in a place where they could be undisturbed, and included local telephone numbers and e-mail addresses to contact if help was desired. Additionally, in Belgium, Estonia and Sweden the participants had the opportunity to complete the questionnaires at the clinic, and measures were taken to avoid accompanying persons being present while they filled out the survey. Formal approvals of local ethical committees and data protection agencies were obtained at all sites, as listed below.

*Belgium:* The Ethical Committee of Ghent University acted as the central ethical committee for the study; U(Z) Gent, 22012008/B67020072813, date of approval: 1<sup>st</sup> February 2008, Waregem hospital date added: 21<sup>st</sup> October 2008. *Iceland:* The scientific board approved the study (24.06.2008-VSN-b2008030024/03-15) according to Icelandic regulations, date: 24<sup>th</sup> June 2008. In *Denmark*, even though ethical approval for non-invasive studies is not required, the study was presented to the Research Ethics Committee of the Capital Region, who found no objections to the study (H-A-2008-002), date: 11<sup>th</sup> February 2008. Permission was obtained from the Danish Data Protection Agency (J.nr. 2007-41-1663). In *Estonia*, ethical permission was given by the Ethics Review Committee on Human Research of the University of Tartu, Estonia; 190/M-29, 192/-22, 196/X-2, date: 17<sup>th</sup> December 2007, East-Tallinn Central Hospital added: 19<sup>th</sup> January 2009, Russian language and prolonged period added: 22<sup>nd</sup> February 2010, East-Viru Central Hospital added: 26<sup>th</sup> April 2010. In *Norway*, the Regional Committee for Medical Research Ethics in North approved the study (72/2006), date: 29<sup>th</sup> August 2007; and the Data Inspectorate (NSD) (15214/3/) also approved the study, date: 19<sup>th</sup> December 2007. In *Sweden*, the study was approved by the Regional Ethical Committee in Stockholm (2006/354-31/1), date: 14<sup>th</sup> June 2006.

The data was anonymised prior to analysis.

#### *Statistical analysis*

Pearson's  $\chi^2$  test was applied to assess demographic and abuse differences between countries. Level of significance was set at  $p < 0.05$ , two sided Kruskal-Wallis test was used to compare medians between countries for the visual analogue scale (VAS) scores for current suffering. The correlation between the level of severity of emotional, physical, and sexual abuse and current suffering was tested by Goodman-Kruskal gamma (G). For comparison between countries of the proportion of lifetime abuse for each type of abuse we calculated

odds ratios (OR) with 95% confidence intervals using logistic regression analysis adjusting for age, education and gestational age when completing the questionnaire with the largest group Norway as a reference. Analyses were done in PASW Statistics version 18.0 (SPSS Inc., Chicago, IL, USA).

## Results

In total of 7200 women responded, 26 women were excluded due to missing response to the NorAQ, leaving a total of 7174 in the study: 861 from Belgium, 602 from Iceland, 1290 from Denmark, 975 from Estonia, 2424 Norway and 1022 from Sweden. Socio-demographic characteristics are presented in Table 1. Significant differences between countries in our sample were observed: nearly a quarter of the women were below 25 years of age in Estonia, while this was so only for around 3 % in Denmark. Norway had the highest proportion of educated women (13 years and more), while the lowest proportion was found in Estonia. Most women were married or cohabiting. Iceland and Estonia had the highest proportion of women not married or cohabiting as well as the highest proportion of women who were unemployed or on social benefit.

In all 3530 women (49.2 %) reported any type of emotional, physical or sexual abuse, 34.8 % when excluding mild physical abuse. Of all the women, 523 (7.3%) reported emotional abuse only, 460 (6.4%) sexual abuse only and 492 (6.9%) physical abuse only (excluding mild physical abuse). One hundred and eighty-eight (2.6%) women reported both emotional and sexual abuse, 355 (4.9%) emotional and physical abuse, 187 (2.6%) physical and sexual abuse, and 294 (4.1%) all three types of abuse.

Tables 2–4 show the proportion of women for each country who reported emotional, physical and sexual abuse by age at time of abuse, severity of the abuse, whether it had occurred within the last year, lifetime abuse, and current suffering. Current moderate or severe suffering from reported emotional abuse was highest among Icelandic women (88.8%) and lowest among Estonian women (68.1%) (Table 2). Seventy percent of the Icelandic women who reported the experience of physical abuse (excluding mild) reported current moderate or severe suffering, compared to 46% of Estonian women (Table 3). The proportion of women reporting no current suffering from their abuse was highest among women who had reported physical abuse, 4.9% (excluding mild physical abuse) compared to 21.3% for emotional abuse and 28.6% for sexual abuse (Tables 2–4). The median scores ranged from 0 for physical abuse only for Denmark, Estonia and Norway to 4 for emotional

and sexual abuse combined for Iceland (Table S2). On the whole women reporting having experienced more than one type of abuse also reported a higher median score for suffering compared to women reporting only one type of abuse, while three types of abuse for the majority of the countries had the highest score (Supporting Information Table S2). The strength of the correlation between severity of each type of abuse and level of suffering was overall moderate ( $G=0.2$ ,  $p<0.001$ ) for emotional abuse and strong for sexual abuse ( $G=0.4$ ,  $p<0.001$ ) and for physical abuse ( $G=0.47$ ,  $p<0.001$ ), although differences existed between countries. For Iceland and Estonia, there was no correlation between degree of suffering and the severity of the emotional abuse.

For all the categories of “any abuse” (excluding mild physical abuse), Estonia had the highest prevalence, with 45.4% reporting any lifetime abuse and 6.5% any current abuse (Table 5). Belgium had the lowest prevalence, 23.3% for any lifetime abuse (excluding mild physical abuse) and 3.0% for any abuse during the past 12 months. Adjusted analyses showed that the adjusted odds for Estonian women to report any lifetime emotional and/or physical abuse (excluding mild physical abuse) was significantly higher compared to Norway, OR 1.63 (1.36–1.95) and 1.54 (1.29–1.84) respectively (Table 6). Belgian and Danish women were significantly less likely to report physical abuse, OR 0.36 (0.28–0.46) and 0.60 (0.49–0.73) respectively; as well as sexual abuse, OR 0.42 (0.32–0.55) and 0.73 (0.60–0.90) respectively. Adjustment had no effect on the significance levels and only marginally altered the odds ratios.

## Discussion

This is the first European multi-country study on the prevalence of different types of abuse among women attending routine antenatal care. Our data suggests that a history of abuse among pregnant women attending routine antenatal care is common. The prevalence of the different types of abuse varied significantly between the participating countries, with or without adjusting for age, education and gestational length at time of participation. The prevalence of current abuse was low. About one in ten women reported severe suffering from the experienced abuse.

In our study women were asked if they had experienced the reported abuse during the past 12 months. Women were on average mid-way through their pregnancy when they filled out the questionnaire. As such we do not report abuse which happened only during pregnancy but current abuse. The prevalence of recent abuse in our study is consistent with 12-month

Accepted Article

estimates from other European settings when measured during pregnancy, such as in Norway (5%) (21), England (1%–5%) (22), Belgium (3.0–3.9%) (23), Sweden ( 2.8%) (24) and Denmark (2.8% during pregnancy) (25). A number of studies report the prevalence of abuse in high income settings separately for the year before pregnancy and during pregnancy, thus complicating comparison to our findings (11, 23, 25). On the whole, the reported prevalence of abuse is higher the year before pregnancy than during pregnancy (7, 11, 18, 23, 26). This is consistent with the protective effect some research claims pregnancy can have, while other studies have noted an increase in particular of emotional and sexual abuse during pregnancy and of the severity and frequency of the abuse (11). Our study did not investigate these aspects of abuse. Alternatively, it may be only the reporting of the abuse which is reduced and not the occurrence.

Our study suggests that in general fewer women suffered from physical abuse and more from emotional abuse, which is consistent with other reports (7). However, it should be noted that 61.5% of those experiencing lifetime emotional abuse (n=1360) were experiencing at least one other kind of abuse as well. Also consistent with other studies we observed that suffering was less when women had reported the experience of only one type of abuse, compared to women reporting two or three types of abuse. In addition, our results showed that the severity of the abuse on the whole corresponded with the degree of current suffering. This suggests a general agreement between researchers and abused women that multiple kinds and severity of abuse is associated with the most suffering. Difference between countries were present which could be due to cultural and contextual differences (27). It may be that in a society with a higher tolerance for violence, the victims tend to regard their experiences as less offensive.

Further, not all pregnant women with a history of abuse report that they suffer from the abuse, or at least not to a great extent. It may be that these women have recovered with or without the help of others, experienced only a single event or a very mild form of abuse only, or had a more resilience. However, we cannot exclude that as a coping mechanism women with abusive experiences might repress their feeling of suffering.

The prevalence of lifetime experience of abuse among pregnant women is relevant for two reasons. Firstly, the past experience may have physical and psychological consequences for the current pregnancy (6). Secondly, women who report previous abuse may be at an increased risk of abuse during pregnancy (11). The lifetime prevalence of any abuse of 23–45% found in our study is consistent with those reported among pregnant women in other European studies, such as 32 % in Norway (21), 23.5% in England (22), 34.5% in Denmark

(25), 19.4% in Sweden (24), 27.6% in Belgium (23). The lifetime prevalence in our study is expected to be lower compared to studies including women of all ages as older women have had a longer lifetime to accumulate abuse in. In our study Estonia had the highest prevalence of any abuse and Belgium the lowest. This could be due to social and cultural differences in what are considered abusive behaviors, which become apparent when abuse is defined by descriptive questions.

Women were recruited while attending routine antenatal care, aiming at an unselected population which would be representative for pregnant women in these countries. While the varying response rate for the participating countries causes concern it is likely that differences in recruitment method played a role. In some of the places women and staff may be frequently asked to participate in research which may reduce their willingness to contribute. The average age of women in the country samples of our study compared well to the average age of pregnant women in the participating countries. Participants in our study had a higher education than the pregnant population in their respective countries 59–72% had more than 13 years of education, compared to national averages of 39–65%. In all participating countries, except for Iceland and Norway, the proportion of nulliparous women was slightly higher among participants (45–54%) than the country average (43–47%). In Belgium the sample was entirely Flemish. In Estonia the proportion of Estonian speaking women (80%) and Russian speaking women (20%) participating in the study are similar to the national proportion of the country.

We used an instrument previously used in a multi-country study (19) but so far only validated in a Swedish population (20, 28). In spite of quality translation into the various languages, the validity may have varied and thus influence the estimates. Using descriptive questions however, is a strength as it allows the researchers to define the abuse and not the participants. Our study was based on self-reported abuse. The results may have been different if personal interviews had been conducted. However, previous studies have found disclosure of sensitive topics higher in self-administered modes compared with face-to-face (29). Obstetricians and midwives meeting women in routine antenatal care should be aware that a high proportion of the women they meet have a history of abuse. Some countries have implemented routine screening to identify current victims of intimate partner violence in antenatal care. It appears that not only is current ongoing abuse of concern but also women with current suffering from earlier abuse could benefit from being identified and receiving specialized care.

## Funding

The Bidens study was supported by the Daphne II Program, European Commission for Freedom, Security, and Justice, Brussels, Belgium (Grant no. JLS/2006/DAP-1/242/W30-CE-0120887/00-87).

## References

1. Garcia-Moreno C. Overview. 2012. In: Understanding and addressing violence against women [Internet].  
<http://www.who.int/reproductivehealth/publications/violence/9789241548595/en/index.html>: WHO.
2. United Nations, General Assembly. Declaration on the elimination of violence against women. Proceedings of the 85th Plenary Meeting, Geneva. 1993.
3. Garcia-Moreno C. Intimate partner violence. 2012. In: Understanding and addressing violence against women [Internet].  
<http://www.who.int/reproductivehealth/publications/violence/9789241548595/en/index.html>: WHO.
4. Ikossi DG, Lazar AA, Morabito D, Fildes J, Knudson MM. Profile of mothers at risk: an analysis of injury and pregnancy loss in 1,195 trauma patients. *J Am Coll Surg*. 2005;200:49-56.
5. Palladino CL, Singh V, Campbell J, Flynn H, Gold KJ. Homicide and suicide during the perinatal period: findings from the National Violent Death Reporting System. *Obstet Gynecol*. 2011;118:1056-63.
6. Ellsberg M, Jansen HA, Heise L, Watts CH, Garcia-Moreno C. Intimate partner violence and women's physical and mental health in the WHO multi-country study on women's health and domestic violence: an observational study. *Lancet*. 2008;371:1165-72.
7. Garcia-Moreno C, Jansen HAFM, Ellsberg M, Heise L, Watts C. WHO Multi-country Study on Women's Health and Domestic Violence against Women Initial results on prevalence, health outcomes and women's responses WHO, editor. Geneva, Switzerland 2005.
8. Campbell JC. Health consequences of intimate partner violence. *Lancet*. 2002;359:1331-6.
9. Howard LM, Oram S, Galley H, Trevillion K, Feder G. Domestic violence and perinatal mental disorders: a systematic review and meta-analysis. *PLoS Med*. 2013;10:e1001452.

- Accepted Article
10. O'Reilly R. Domestic violence against women in their childbearing years: a review of the literature. *Contemp Nurse*. 2007;25:13-21.
  11. Taillieu TL, Brownridge DA. Violence against pregnant women: Prevalence, patterns, risk factors, theories, and directions for future research. *Aggr Violent Behav*. 2010;15:14-35.
  12. Garcia-Moreno C. Intimate-partner violence and fetal loss. *Lancet*. 2009;373:278-9.
  13. Leone JM, Lane SD, Koumans EH, DeMott K, Wojtowycz MA, Jensen J, et al. Effects of intimate partner violence on pregnancy trauma and placental abruption. *J Womens Health (Larchmt)*. 2010;19:1501-9.
  14. Rodrigues T, Rocha L, Barros H. Physical abuse during pregnancy and preterm delivery. *Am J Obstet Gynecol*. 2008;198:171 e1-6.
  15. Kendall-Tackett K. The health effects of childhood abuse: four pathways by which abuse can influence health. *Child Abuse Negl*. 2002;26:715-29.
  16. Leeners B, Stiller R, Block E, Gorres G, Rath W. Pregnancy complications in women with childhood sexual abuse experiences. *J Psychosom Res*. 2010;69:503-10.
  17. Chambliss LR. Intimate partner violence and its implication for pregnancy. *Clin Obstet Gynecol*. 2008;51:385-97.
  18. Devries KM, Kishor S, Johnson H, Stöckl H, Bacchus LJ, Garcia-Moreno C, et al. Intimate partner violence during pregnancy: analysis of prevalence data from 19 countries. *Reprod Health Matters*. 2010;18:158-70.
  19. Wijma B, Schei B, Swahnberg K, Hilden M, Offerdal K, Pikarinen U, et al. Emotional, physical, and sexual abuse in patients visiting gynaecology clinics: a Nordic cross-sectional study. *Lancet*. 2003;361:2107-13.
  20. Swahnberg IM, Wijma B. The NorVold Abuse Questionnaire (NorAQ): validation of new measures of emotional, physical, and sexual abuse, and abuse in the health care system among women. *Eur J Public Health*. 2003;13:361-6.
  21. Sorbo MF, Grimstad H, Bjørngaard JH, Schei B, Lukasse M. Prevalence of sexual, physical and emotional abuse in the Norwegian mother and child cohort study. *BMC Public Health*. 2013;13:186.
  22. Bacchus L, Mezey G, Bewley S. Domestic violence: prevalence in pregnant women and associations with physical and psychological health. *Eur J Obstet Gynecol Reprod Biol*. 2004;113:6-11.
  23. Roelens K, Verstraelen H, Van Egmond K, Temmerman M. Disclosure and health-seeking behaviour following intimate partner violence before and during pregnancy in

Flanders, Belgium: a survey surveillance study. *Eur J Obstet Gynecol Reprod Biol.* 2008;137:37-42.

24. Stenson K, Heimer G, Lundh C, Nordström ML, Saarinen H, Wenker A. The prevalence of violence investigated in a pregnant population in Sweden. *J Psychosom Obstet Gynaecol.* 2001;22:189-97.

25. Finnbogadóttir H, Dejin-Karlsson E, Dykes AK. A multi-centre cohort study shows no association between experienced violence and labour dystocia in nulliparous women at term. *BMC Pregnancy Childbirth.* 2011;11:14.

26. Stenson K, Heimer G, Lundh C, Nordström ML, Saarinen H, Wenker A. Lifetime prevalence of sexual abuse in a Swedish pregnant population. *Acta Obstet Gynecol Scand.* 2003;82:529-36.

27. Lindhorst T, Tajima E. Reconceptualizing and operationalizing context in survey research on intimate partner violence. *J Interpers Violence.* 2008;23:362-88.

28. Nybergh L, Taft C, Krantz G. Psychometric properties of the WHO Violence Against Women instrument in a female population-based sample in Sweden: a cross-sectional survey. *BMJ Open.* 2013;3: pii: e002053.

29. Bowling A. Mode of questionnaire administration can have serious effects on data quality. *J Public Health.* 2005;27:281-91.

Table 1. Baseline characteristics for pregnant women in the Bidens cohort study, 2008–2010.

	<b>Belgium</b> n=861 n (%)	<b>Iceland</b> n=602 n (%)	<b>Denmark</b> n=1290 n (%)	<b>Estonia</b> n=975 n (%)	<b>Norway</b> n=2424 n (%)	<b>Sweden</b> n=1022 n (%)	<b>Total</b> N=7174 n (%)
<b>Age (years)<sup>†</sup></b>							
< 25 years	149 (17.4)	91 (15.3)	41 (3.2)	238 (24.6)	286 (11.8)	112 (11.1)	917 (12.9)
25 – 34 years	632 (73.9)	396 (66.7)	923 (72.4)	619 (63.9)	1609 (66.5)	708 (70.0)	4887 (68.6)
≥35 years	74 (8.7)	107 (18.0)	311 (24.4)	112 (11.6)	526 (21.7)	191 (18.9)	1321 (18.5)
<b>Education (years attained)<sup>†</sup></b>							
< 9 years	13 (1.5)	44 (7.4)	19 (1.5)	76 (7.9)	58 (2.4)	34 (3.4)	244 (3.5)
10 – 13 years	322 (37.8)	142 (24.0)	119 (9.4)	324 (33.6)	618 (25.8)	307 (30.8)	1832 (25.9)
> 13 years	516 (60.6)	405 (68.5)	1133 (89.1)	564 (58.5)	1723 (71.8)	655 (65.8)	4996 (70.6)
<b>Civil status<sup>‡</sup></b>							
Married/ cohabiting	822 (97.0)	549 (93.1)	1218 (96.0)	913 (94.4)	2314 (96.4)	971 (96.2)	6787 (95.8)
Others	25 (3.0)	41 (6.9)	51 (4.0)	54 (5.6)	86 (3.6)	38 (3.8)	295 (4.2)
<b>Occupation<sup>†</sup></b>							
Employed/ student	604 (71.1)	532 (89.6)	1200 (94.1)	628 (64.8)	2208 (91.6)	926 (91.8)	6098 (85.8)
Pregnancy leave	182 (21.4)	1 (0.2)	19 (1.5)	199 (20.5)	63 (2.6)	23 (2.3)	487 (6.9)
Housewife	19 (2.2)	17 (2.9)	10 (0.8)	83 (8.6)	68 (2.8)	10 (1.0)	207 (2.9)
Unemployed/ social benefits	45 (5.3)	44 (7.4)	46 (3.6)	59 (6.1)	71 (2.9)	50 (5.0)	315 (4.4)

<sup>†</sup>  $p < 0.001$ , <sup>‡</sup>  $p = 0.001$ , Pearson's  $\chi^2$ -test

Table 2. Prevalence of emotional abuse and current suffering among pregnant women in the Bidens cohort study, 2008–2010.

	Belgium n=861 n (%)	Iceland n=602 n (%)	Denmark n=1290 n (%)	Estonia n=975 n (%)	Norway n=2424 n (%)	Sweden n=1022 n (%)	Total N=7174 n (%)
<b>Severity of abuse</b>							
<b>Mild<sup>†</sup></b>							
< 18 years	31 (3.6)	21 (3.5)	66 (5.1)	60 (6.2)	70 (2.9)	27 (2.6)	275 (3.8)
≥ 18 years	19 (2.2)	6 (1.0)	46 (3.6)	21 (2.2)	48 (2.0)	25 (2.4)	165 (2.3)
Both	1 (0.1)	0	17 (1.3)	12 (1.2)	14 (0.6)	10 (1.0)	54 (0.8)
<b>Moderate<sup>‡</sup></b>							
< 18 years	13 (1.5)	13 (2.2)	20 (1.6)	48 (4.9)	35 (1.4)	12 (1.2)	141 (2.0)
≥ 18 years	10 (1.2)	22 (3.7)	12 (0.9)	18 (1.9)	56 (2.3)	23 (2.3)	141 (2.0)
Both	8 (0.9)	7 (1.2)	1 (0.1)	6 (0.6)	8 (0.3)	5 (0.5)	35 (0.5)
<b>Severe<sup>‡</sup></b>							
< 18 years	25 (2.9)	16 (2.7)	36 (2.8)	55 (5.7)	90 (3.7)	28 (2.8)	250 (3.5)
≥ 18 years	28 (3.3)	10 (1.7)	49 (3.8)	43 (4.4)	92 (3.8)	22 (2.2)	244 (3.4)
Both	4 (0.5)	3 (0.5)	5 (0.4)	6 (0.6)	23 (1.0)	14 (1.4)	55 (0.8)
<b>Abuse past 12 months<sup>†</sup></b>							
Any lifetime abuse <sup>‡</sup>	23 (2.7)	11 (1.8)	28 (2.2)	49 (5.0)	57 (2.4)	22 (2.2)	190 (2.7)
Current suffering*	139 (16.2) (%) <sup>*</sup>	98 (16.3) (%) <sup>*</sup>	252 (19.6) (%) <sup>*</sup>	269 (27.7) (%) <sup>*</sup>	436 (18.0) (%) <sup>*</sup>	166 (16.3) (%) <sup>*</sup>	1360 (19.0) (%) <sup>*</sup>
None	29 (20.9)	11 (11.2)	42 (16.7)	80 (29.7)	100 (22.9)	28 (16.9)	290 (21.3)
Moderate	91(65.5)	64 (65.3)	168 (66.7)	157 (58.4)	278 (63.8)	105 (63.3)	863 (63.5)
Severe	13 (9.4 )	23 (23.5)	38 (15.1)	26 (9.7)	50 (11.5)	30 (18.1)	180 (13.2)
Missing	6 (4.3)	0	4 (1.6)	6 (2.2)	8 (1.8)	3 (1.8)	27 (2.0)

<sup>†</sup> NS  $p=0.13$ , <sup>‡</sup> $p<0.001$ , Pearson's  $\chi^2$ -test, \* percentage among women reporting any lifetime abuse

Table 3. Prevalence of physical abuse and current suffering among pregnant women in the Bidens cohort study, 2008–2010.

	Belgium n=861 n (%)	Iceland n=602 n (%)	Denmark n=1290 n (%)	Estonia n=975 n (%)	Norway n=2424 n (%)	Sweden n=1022 n (%)	Total N=7174 n (%)
<b>Severity of abuse</b>							
<b>Mild<sup>†</sup></b>							
< 18 years	105 (12.2)	46 (7.6)	315 (24.4)	93 (9.5)	440 (18.1)	115 (11.3)	1114 (15.3)
≥ 18 years	12 (1.4)	21 (3.5)	82 (6.4)	56 (5.7)	98 (4.0)	43 (4.2)	312 (4.4)
Both	6 (0.7)	6 (1.0)	39 (3.0)	6 (0.6)	59 (2.4)	18 (1.8)	134 (1.9)
<b>Moderate<sup>†</sup></b>							
< 18 years	23 (2.7)	54 (9.0)	46 (3.6)	108 (11.1)	152 (6.3)	40 (3.9)	423 (5.9)
≥ 18 years	15 (1.7)	27 (4.5)	36 (2.8)	56 (5.7)	129 (5.3)	33 (3.2)	296 (4.1)
Both	6 (0.7)	10 (1.7)	2 (0.2)	16 (1.6)	21 (0.9)	12 (1.2)	67 (0.9)
<b>Severe<sup>†</sup></b>							
< 18 years	15 (1.8)	9 (1.5)	19 (1.5)	49 (5.1)	66 (2.7)	15 (1.5)	173 (2.4)
≥ 18 years	24 (2.8)	16 (2.7)	59 (4.6)	66 (6.8)	137 (5.7)	39 (3.8)	341 (4.8)
Both	0	0	1 (0.1)	5 (0.5)	15 (0.6)	7 (0.7)	28 (0.4)
<b>Abuse past 12 months<sup>†</sup></b>							
Any lifetime abuse <sup>†</sup>	11 (1.3) (24.0)	13 (2.2) (31.4)	32 (2.5) (46.5)	27 (2.8) (46.7)	55 (2.3) (46.1)	18 (1.8) (31.5)	156 (2.2) (40.3)
Current suffering <sup>†*</sup>	(%)*	(%)*	(%)*	(%)*	(%)*	(%)*	(%)*
None	106 (51.5)	73 (38.6)	342 (57.1)	266 (58.5)	697 (67.4)	151 (46.9)	1635 (55.6)
Moderate	71 (34.5)	87 (46.0)	178 (29.7)	167 (36.7)	323 (28.9)	135 (41.9)	961 (33.3)
Severe	4 (1.9)	21 (11.1)	16 (2.7)	13 (2.9)	23 (2.1)	24 (7.5)	101 (3.5)
Missing	25 (12.1)	8 (4.2)	63 (10.5)	9 (2.0)	74 (6.6)	12 (3.7)	191 (6.6)
<b>Physical abuse, mild physical abuse excluded</b>							
Any lifetime abuse <sup>†</sup>	83 (9.7)	116 (19.3)	163 (12.6)	300 (30.8)	520 (21.5)	146 (14.3)	1328 (18.5)
Current suffering <sup>†*</sup>	(%)*	(%)*	(%)*	(%)*	(%)*	(%)*	(%)*
None	26 (31.3)	34 (29.3)	66 (40.5)	158 (52.7)	264 (50.8)	48 (32.9)	596 (44.9)
Moderate	53 (63.9)	63 (54.3)	83 (50.9)	127 (42.3)	225 (43.3)	75 (51.4)	626 (47.1)
Severe	4 (4.8)	18 (15.5)	13 (8.0)	12 (4.0)	21 (4.0)	21 (14.4)	89 (6.7)
Missing	0	1 (0.9)	1 (0.6)	3 (1.0)	10 (1.9)	2 (1.4)	17 (1.3)

<sup>†</sup>  $p < 0.001$ , Pearson's  $\chi^2$ -test; \* percentage among women reporting any lifetime abuse

Table 4. Prevalence of sexual abuse and current suffering among pregnant women in the Bidens cohort study, 2008–2010.

	Belgium n=861 n (%)	Iceland n=602 n (%)	Denmark n=1290 n (%)	Estonia n=975 n (%)	Norway n=2424 n (%)	Sweden n=1022 n (%)	Total N=7174 n (%)
<b>Severity of abuse</b>							
<b>Mild<sup>¶</sup></b>							
< 18 years	10 (1.2)	17 (2.8)	38 (2.9)	19 (2.0)	58 (2.4)	20 (2.0)	162 (2.3)
≥ 18 years	1 (0.1)	4 (0.7)	25 (1.9)	9 (0.9)	27 (1.1)	15 (1.5)	81 (1.1)
Both	1 (0.1)	2 (0.3)	2 (0.2)	1 (0.1)	3 (0.1)	0	9 (0.1)
<b>Moderate<sup>§</sup></b>							
< 18 years	21 (2.4)	32 (5.3)	33 (2.6)	56 (5.7)	111 (4.6)	35 (3.4)	288 (4.0)
≥ 18 years	2 (0.2)	4 (0.7)	11 (0.9)	9 (0.9)	27 (1.1)	12 (1.2)	65 (0.9)
Both	0	1 (0.2)	1 (0.1)	1 (0.1)	3 (0.1)	2 (0.2)	8 (0.1)
<b>Severe<sup>†</sup></b>							
< 18 years	23 (2.7)	39 (6.5)	25 (1.9)	43 (4.4)	89 (3.7)	31 (3.0)	250 (3.5)
≥ 18 years	13 (1.5)	21 (3.5)	29 (2.2)	35 (3.6)	94 (3.9)	37 (3.6)	229 (3.2)
Both	0	7 (1.2)	4 (0.3)	2 (0.2)	18 (0.7)	5 (0.5)	36 (0.5)
<b>Abuse past 12 months<sup>  </sup></b>							
	0	3 (0.5)	3 (0.2)	8 (0.8)	12 (0.5)	3 (0.3)	29 (0.4)
<b>Any lifetime abuse<sup>‡</sup></b>							
	71 (8.3)	127 (21.1)	168 (13.0)	175 (18.0)	430 (17.7)	157 (15.5)	1129 (15.7)
<b>Current suffering<sup>‡,*</sup></b>							
	(%)*	(%)*	(%)*	(%)*	(%)*	(%)*	(%)*
None	14 (19.6)	22 (17.3)	52 (31.0)	58 (33.1)	135 (31.4)	42 (26.6)	323 (28.6)
Moderate	48 (67.6)	80 (63.0)	98 (58.3)	99 (56.6)	249 (57.9)	87 (55.1)	661 (58.5)
Severe	3 (4.2)	19 (15.0)	14 (8.3)	13 (7.4)	36 (8.4)	22 (13.9)	107 (9.5)
Missing	6 (8.5)	6 (4.7)	4 (2.4)	5 (2.9)	10 (2.3)	7 (4.4)	38 (3.4)

<sup>¶</sup>  $p=0.01$ , <sup>§</sup>  $p=0.001$ , <sup>†</sup>  $p<0.001$ , <sup>||</sup> NS=0.40, <sup>‡</sup>  $p=0.01$ , Pearson's  $\chi^2$ -test; \* percentage among women reporting any lifetime abuse

Table 5. Prevalence of any childhood, adult, lifetime and current abuse among pregnant women in the Bidens cohort study\*, 2008–2010.

	Belgium	Iceland	Denmark	Estonia	Norway	Sweden	Total
	n=861 n (%)	n=602 n (%)	n=1290 n (%)	n=975 n (%)	n=2424 n (%)	n=1022 n (%)	N=7174 n (%)
<b>Any childhood abuse<sup>†</sup></b>	139 (16.1)	171 (28.4)	255 (19.8)	337 (34.6)	583 (24.1)	208 (20.4)	1693 (23.6)
<b>Any adult abuse<sup>†</sup></b>	100 (11.6)	99 (16.4)	259 (20.1)	217 (22.3)	541 (22.3)	197 (19.3)	1413 (19.7)
<b>Any lifetime abuse<sup>†</sup></b>	200 (23.2)	214 (35.5)	433 (33.6)	443 (45.4)	900 (37.1)	309 (30.2)	2499 (34.8)
<b>Any abuse past 12 months<sup>†</sup></b>	26 (3.0)	20 (3.3)	42 (3.3)	63 (6.5)	89 (3.7)	31 (3.0)	271 (3.8)

\* Women with only mild physical abuse were excluded in these analyses

<sup>†</sup>  $p < 0.001$ , Pearson's  $\chi^2$ -test

Table 6. Crude and adjusted\* odds ratio (OR) (95 % confidence interval) of emotional, physical and sexual lifetime abuse, among pregnant women in the Bidens cohort study, 2008–2010.

Country	Emotional abuse	Physical abuse	Physical abuse, excluding mild physical abuse	Sexual abuse
	Adjusted OR	Adjusted OR	Adjusted OR	Adjusted OR
<b>Norway</b>	1	1	1	1
<b>Belgium</b>	0.87 (0.71 – 1.08)	0.36 (0.30 - 0.43)	0.36 (0.28 - 0.46)	0.42 (0.32 - 0.55)
<b>Iceland</b>	0.84 (0.66 - 1.08)	0.51 (0.42 - 0.62)	0.83 (0.65 - 1.05)	1.16 (0.92 - 1.47)
<b>Denmark</b>	1.19 (1.00 - 1.43)	1.07 (0.93 - 1.23)	0.60 (0.49 - 0.73)	0.73 (0.60 - 0.90)
<b>Estonia</b>	1.63 (1.36 - 1.95)	0.99 (0.85 - 1.16)	1.54 (1.29 - 1.84)	0.99 (0.81 - 1.22)
<b>Sweden</b>	0.84 (0.68 - 1.03)	0.52 (0.44 - 0.61)	0.55 (0.44 - 0.68)	0.81 (0.65 - 0.99)

\*Adjusted for age, education and gestational week for filling out the questionnaire.